

## References

- [Alcalà-Quintana and García-Perez(2004)] Alcalà-Quintana, R., García-Perez, M., 2004. The role of parametric assumptions in adaptive bayesian estimation. *Psych. Meth.* 9 (2), 250–271.
- [Anderson et al.(1946)Anderson, McCarthy, and Tukey] Anderson, T., McCarthy, P., Tukey, J., 1946. staircase method of sensitivity testing. Naval Ordinance Report 65-46, Statistical Research Group, Princeton University, Princeton, NJ, USA.
- [ASTM(1991)] ASTM, 1991. Standard test method for estimating acute oral toxicity in rats. American Society for Testing and Materials, Philadelphia, USA, designation E 1163-90.
- [Auerswald et al.(2005)Auerswald, Pfeiffer, Behrends, Burkhardt, and Olthoff] Auerswald, K., Pfeiffer, F., Behrends, K., Burkhardt, U., Olthoff, D., 2005. Pain on injection with propofol. *ANASTH. INTENSIV. NOTFAL. SCHMERZ.* 40 (5), 259–266.
- [Babb et al.(1998)Babb, Rogatko, and Zacks] Babb, J., Rogatko, A., Zacks, S., 1998. Cancer phase I clinical trials: Efficient dose escalation with overdose control. *Stat. Med.* 17, 1103–1120.
- [Banerjee and Wellner(2005)] Banerjee, M., Wellner, J. A., 2005. Score statistics for current status data: comparisons with likelihood ratio and Wald statistics. *Int. J. Biostat.* 1, Art. 3, 29 pp. (electronic).
- [Barlow et al.(1972)Barlow, Bartholomew, Bremner, and Brunk] Barlow, R. E., Bartholomew, D. J., Bremner, J. M., Brunk, H. D., 1972. *Statistical Inference Under Order Restrictions. The Theory and Application of Isotonic Regression.* John Wiley & Sons, London-New York-Sydney.
- [Bhananker et al.(2007)Bhananker, Tontisirin, Oron, Muangman, and Souter] Bhananker, S. M., Tontisirin, N., Oron, A. P., Muangman, S. L., Souter, M. J., 2007. Optimal ratio of thiopental-propofol mixture to eliminate pain on injection of propofol. Poster Session.
- [Bortot and Giovagnoli(2005)] Bortot, P., Giovagnoli, A., 2005. Up-and-down experiments of first and second order. *J. Statist. Plann. Inference* 134 (1), 236–253.
- [Braam and van der Zwaag(1998)] Braam, J., van der Zwaag, S., 1998. A statistical evaluation of the staircase and the ArcSin root P methods for determining the fatigue limit. *J. Testing Eval.* 26, 125–131.

- [Brownlee et al.(1953)Brownlee, Hodges, and Rosenblatt] Brownlee, K. A., Hodges, Jr., J. L., Rosenblatt, M., 1953. The up-and-down method with small samples. *J. Amer. Statist. Assoc.* 48, 262–277.
- [Buoen et al.(2005)Buoen, Bjerrum, and Thomsen] Buoen, C., Bjerrum, O. J., Thomsen, M. S., 2005. How first-time-in-human studies are being performed: a survey of phase i dose-escalation trials in healthy volunteers published between 1995 and 2004. *J. Clin. Pharmacol.* 45, 1123–1136.
- [Camorcia et al.(2004)Camorcia, Capogna, Lyons, and Columb] Camorcia, M., Capogna, G., Lyons, G., Columb, M., 2004. Epidural test dose with levobupivacaine and ropivacaine: determination of ED50 motor block after spinal administration. *Brit. J. Anaesthes* 92 (6), 850–853.
- [Capogna et al.(2001)Capogna, Parpaglioni, Lyons, Columb, and Celleno] Capogna, G., Parpaglioni, R., Lyons, G., Columb, M., Celleno, D., 2001. Minimum analgesic dose of epidural sufentanil for first-stage labor analgesia. *Anesthesiology* 94, 740–744.
- [Chao and Fuh(2001)] Chao, M. T., Fuh, C. D., 2001. Bootstrap methods for the up and down test on pyrotechnics sensitivity analysis. *Statist. Sinica* 11 (1), 1–21.
- [Chao and Fuh(2003)] Chao, M. T., Fuh, C. D., 2003. Why wild values occur in pyrotechnic sensitivity analysis. *Propell. Explos. Pyrotech.* 28 (4), 216–218.
- [Cheung and Chappel(2002)] Cheung, Y., Chappel, R., 2002. A simple technique to evaluate model sensitivity in the continual reassessment method. *Biometrics* 58 (3), 671–674.
- [Chevret(2006)] Chevret, S. E., 2006. *Statistical Methods for Dose-finding Experiments*. Statistics in Practice. John Wiley & Sons, Chichester.
- [Choi(1971)] Choi, S., 1971. An investigation of Wetherill’s method of estimation for the up-and-down experiment. *Biometrics* 27 (4), 961–970.
- [Choi(1990)] Choi, S., 1990. Interval estimation of the LD50 based on an up-and-down experiment. *Biometrics* 46 (2), 485–492.
- [Chung(1960)] Chung, K., 1960. *Statistical Inference for Markov Chains*. Springer, Berlin.
- [Davison and Hinkley(1995)] Davison, A., Hinkley, D., 1995. *Bootstrap Methods and their Applications*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, Cambridge.

- [Derman(1957)] Derman, C., 1957. Non-parametric up-and-down experimentation. *Ann. Math. Statist.* 28, 795–798.
- [Diaconis and Stroock(1991)] Diaconis, P., Stroock, D., 1991. Geometric bounds for eigenvalues of Markov chains. *Ann. Appl. Probab.* 1 (1), 36–61.
- [Dilleen et al.(2003)] Dilleen, M., Heimann, G., Hirsch, I., 2003. Non-parametric estimators of monotonic dose-response curve and bootstrap intervals. *Stat. Med.* 22, 869–882.
- [Dixon(1965)] Dixon, W. J., 1965. The up-and-down method for small samples. *J. Amer. Statist. Assoc.* 60, 967–978.
- [Dixon and Mood(1948)] Dixon, W. J., Mood, A., 1948. A method for obtaining and analyzing sensitivity data. *J. Amer. Statist. Assoc.* 43, 109–126.
- [Drover et al.(2004)] Drover, D., Litalien, C., Wellis, V., Shafer, S., Hammer, G., 2004. Determination of the pharmacodynamic interaction of propofol and remifentanyl during esophagogastroduodenoscopy in children. *Anesthesiology* 100, 1382–1386.
- [Durham and Flournoy(1994)] Durham, S. D., Flournoy, N., 1994. Random walks for quantile estimation. In: *Statistical decision theory and related topics, V* (West Lafayette, IN, 1992). Springer, New York, pp. 467–476.
- [Durham and Flournoy(1995)] Durham, S. D., Flournoy, N., 1995. Up-and-down designs. I. Stationary treatment distributions. In: *Adaptive designs* (South Hadley, MA, 1992). Vol. 25 of *IMS Lecture Notes Monogr. Ser. Inst. Math. Statist.*, Hayward, CA, pp. 139–157.
- [Durham et al.(1995)] Durham, S. D., Flournoy, N., Montazer-Haghighi, A. A., 1995. Up-and-down designs. II. Exact treatment moments. In: *Adaptive designs* (South Hadley, MA, 1992). Vol. 25 of *IMS Lecture Notes Monogr. Ser. Inst. Math. Statist.*, Hayward, CA, pp. 158–178.
- [Durham et al.(1997)] Durham, S. D., Flournoy, N., Rosenberger, W., 1997. A random walk rule for phase I clinical trials. *Biometrics* 53 (2), 745–760.
- [Efron and Tibshirani(1993)] Efron, B., Tibshirani, R. J., 1993. An introduction to the bootstrap. Vol. 57 of *Monographs on Statistics and Applied Probability*. Chapman and Hall, New York.
- [Fill(1991)] Fill, J. A., 1991. Eigenvalue bounds on convergence to stationarity for nonreversible Markov chains, with an application to the exclusion process. *Ann. Appl. Probab.* 1 (1), 62–87.

- [Fisher(2007)] Fisher, D., 2007. What if half of your patients moved (or remembered or did something else bad) at incision? *Anesthesiology* 107 (1), 1–2, editorial View.
- [Friedman and Tibshirani(1984)] Friedman, J., Tibshirani, R., 1984. The monotone smoothing of scatterplots. *Technometrics* 26, 243–250.
- [García-Perez(1998)] García-Perez, M., 1998. Forced-choice staircases with fixed step sizes: asymptotic and small-sample properties. *Vision Res.* 38, 1861–1881.
- [García-Perez and Alcalà-Quintana(2005)] García-Perez, M., Alcalà-Quintana, R., 2005. Sampling plans for fitting the psychometric function. *Span. J. Psych.* 8 (2), 256–289.
- [Garrett-Mayer(2006)] Garrett-Mayer, E., 2006. The continual reassessment method for dose-finding studies: a tutorial. *Clinical Trials* 3 (1), 57–71.  
URL <http://ctj.sagepub.com/cgi/content/abstract/3/1/57>
- [Gasparini and Eisele(2000)] Gasparini, M., Eisele, J., 2000. A curve-free method for phase I clinical trials. *Biometrics* 56, 609–615.
- [Gezmu(1996)] Gezmu, M., 1996. The geometric up-and-down design for allocating dosage levels. Ph.D. thesis, American University, Washington, DC, USA.
- [Gezmu and Flournoy(2006)] Gezmu, M., Flournoy, N., 2006. Group up-and-down designs for dose-finding. *J. Statist. Plann. Inference* 136 (6), 1749–1764.
- [Giovagnoli(2004)] Giovagnoli, A., 2004. Asymptotic properties of biased coin designs for treatment allocation. In: *mODa 7—Advances in model-oriented design and analysis*. *Contrib. Statist. Physica*, Heidelberg, pp. 81–88.
- [Giovagnoli and Pintacuda(1998)] Giovagnoli, A., Pintacuda, N., 1998. Properties of frequency distributions induced by general “up-and-down” methods for estimating quantiles. *J. Statist. Plann. Inference* 74 (1), 51–63.
- [Goodman et al.(1995)Goodman, Zahurak, and Piantadosi] Goodman, S., Zahurak, M., Piantadosi, S., 1995. Some practical improvements in the continual reassessment method for phase I studies. *Stat. Med.* 14, 1149–1161.
- [Haines et al.(2003)Haines, Perevozskaya, and Rosenberger] Haines, L., Perevozskaya, I., Rosenberger, W., 2003. Bayesian optimal designs for phase I clinical trials. *Biometrics* 59 (3), 591–600.

- [Hastie et al.(2001)Hastie, Tibshirani, and Friedman] Hastie, T., Tibshirani, R., Friedman, J., 2001. *The Elements of Statistical Learning*. Springer Series in Statistics. Springer-Verlag, New York, data mining, inference, and prediction.
- [Hastie and Tibshirani(1990)] Hastie, T. J., Tibshirani, R. J., 1990. *Generalized Additive Models*. Vol. 43 of *Monographs on Statistics and Applied Probability*. Chapman and Hall Ltd., London.
- [He and Shi(1998)] He, X., Shi, P., 1998. Monotone B-spline smoothing. *J. Amer. Statist. Assoc.* 93, 643–650.
- [Hughes(1995)] Hughes, B. D., 1995. *Random Walks and Random Environments*. Vol. 1. Oxford Science Publications. The Clarendon Press Oxford University Press, New York, random walks.
- [Ivanova(2006)] Ivanova, A., 2006. Escalation, up-and-down and A+B designs for dose-finding trials. *Stat. Med.* 25, 3668–3678.
- [Ivanova and Flournoy(2006)] Ivanova, A., Flournoy, N., 2006. Up-and-down designs in toxicity studies. In: *Statistical Methods for Dose-finding Experiments*. *Statistics in Practice*. John Wiley & Sons, Chichester, pp. 115–130, ed: S. Chevret.
- [Ivanova et al.(2007)Ivanova, Flournoy, and Chung] Ivanova, A., Flournoy, N., Chung, Y., 2007. Cumulative cohort design for dose-finding. *J. Statist. Plann. Inference* 137, 2316–2327.
- [Ivanova et al.(2003)Ivanova, Haghghi, Mohanty, and Durham] Ivanova, A., Haghghi, A., Mohanty, S., Durham, S., 2003. Improved up-and-down designs for phase I trials. *Stat. Med.* 22, 69–82.
- [JSME(1981)] JSME, 1981. Standard method of statistical fatigue testing. Japan Society of Mechanical Engineers, Japan, jSME S 002.
- [Kershaw(1985)] Kershaw, C., 1985. Asymptotic properties of  $w$ , an estimator of the  $ed_{50}$  suggested for use in up-and-down experiments in bioassay. *Ann. Statist.* 13 (2), 85–94.
- [Kershaw(1987)] Kershaw, C., 1987. A comparison of the estimators of the  $ED_{50}$  in up-and-down experiments. *J. Stat. Comp. Simul.* 27.
- [King-Smith et al.(1994)King-Smith, Grigsby, Vingrys, Benes, and Supowit] King-Smith, P., Grigsby, S., Vingrys, A., Benes, S., Supowit, A., 1994. Efficient and unbiased modifications of the QUEST threshold method: theory, simulations, experimental evaluation and practical implementation. *Vision Res.* 34, 885–912.

- [Komori and Hirose(2000)] Komori, Y., Hirose, H., 2000. An easy parameter estimation by the EM algorithm in the new up-and-down method. *IEEE Trans. Dielect. Elect. Insul.* 7 (6), 838–842.
- [Korn et al.(1994)] Korn, Midthune, Chen, Rubinstein, Christian, and Simon] Korn, E., Midthune, D., Chen, T., Rubinstein, L., Christian, M., Simon, R., 1994. A comparison of 2 phase-I trial designs. *Stat. Med.* 13 (18), 1799–1806.
- [Lagoda and Sonsino(2004)] Lagoda, T., Sonsino, C., 2004. Comparison of different methods for presenting variable amplitude loading fatigue results. *Mat. Wissen. Werkstoff* 35 (1), 13–20.
- [Lai(2001)] Lai, T. L., 2001. Sequential analysis: some classical problems and new challenges. *Statist. Sinica* 11 (2), 303–408, with comments and a rejoinder by the author.
- [Lai(2003)] Lai, T. L., 2003. Stochastic approximation. *Ann. Statist.* 31 (2), 391–406, dedicated to the memory of Herbert E. Robbins.
- [Lai(2004)] Lai, T. L., 2004. Interim and terminal analyses of clinical trials with failure-time endpoints and related group sequential designs. In: *Applied sequential methodologies*. Vol. 173 of *Statist. Textbooks Monogr.* Dekker, New York, pp. 193–218.
- [Lai and Shih(2004)] Lai, T. L., Shih, M.-C., 2004. Power, sample size and adaptation considerations in the design of group sequential clinical trials. *Biometrika* 91 (3), 507–528.
- [Leung and Wang(2001)] Leung, D. H.-Y., Wang, Y.-G., 2001. Isotonic designs for phase I trials. *Cont. Clin. Trials* 22 (2), 126–138.
- [Lichtman(1998)] Lichtman, A., 1998. The up-and-down method substantially reduces the number of animals required to determine antinociceptive ED50 values. *J. Pharma. and Toxic. Meth.* 40 (2), 81–85.
- [Lin et al.(2001)] Lin, Lee, and Lu] Lin, S., Lee, Y., Lu, M., 2001. Evaluation of the staircase and the accelerated test methods for fatigue limit distributions. *Int. J. Fat.* 23.
- [Lin and Shih(2001)] Lin, Y., Shih, W., 2001. Statistical properties of the traditional algorithm-based designs for phase I cancer clinical trials. *Biostatistics* 2 (2), 203–215.
- [Mammen(1991)] Mammen, ., 1991. Estimating a smooth monotone regression function. *Ann. Statist.* 19, 724–740.

- [Marvit et al.(2003)Marvit, Florentine, and Buus] Marvit, P., Florentine, M., Buus, S., 2003. A comparison of psychophysical procedures for level-discrimination thresholds. *J. Acoust. Soc. Am.* 113 (6), 3348–3361.
- [Mathew et al.(2004)Mathew, Thall, Jones, Perez, Bucana, Troncoso, Kim, Fidler, and Logothetis] Mathew, P., Thall, P. F., Jones, D., Perez, C., Bucana, C., Troncoso, P., Kim, S.-J., Fidler, I. J., Logothetis, C., 2004. Platelet-Derived Growth Factor Receptor Inhibitor Imatinib Mesylate and Docetaxel: A Modular Phase I Trial in Androgen-Independent Prostate Cancer. *J Clin Oncol* 22 (16), 3323–3329.  
URL <http://www.jco.org/cgi/content/abstract/22/16/3323>
- [NIEHS(2001)] NIEHS, 2001. The Revised Up-and-Down Procedure: A Test Method for Determining the Acute Oral Toxicity of Chemicals. National Institute of Environmental Health Sciences, National Institute of Health, Washington D.C., USA, NIH publication No. 02-4501.
- [OECD(1998)] OECD, 1998. The Revised Up-and-Down Procedure: A Test Method for Determining the Acute Oral Toxicity of Chemicals. Organisation for Economic Co-operation and Development, Paris, France.
- [O’Quigley(2002)] O’Quigley, J., 2002. Curve-free and model-based continual reassessment designs. *Biometrics* 58, 245–249.
- [O’Quigley and Chevret(1991)] O’Quigley, J., Chevret, S., 1991. Methods for dose finding studies in cancer clinical-trials - a review and results of a monte-carlo study. *Stat. Med.* 10 (11), 1647–1664.
- [O’Quigley et al.(1990)O’Quigley, Pepe, and Fisher] O’Quigley, J., Pepe, M., Fisher, L., 1990. Continual reassessment method: a practical design for phase 1 clinical trials in cancer. *Biometrics* 46 (1), 33–48.
- [O’Quigley and Zohar(2006)] O’Quigley, J., Zohar, S., 2006. Experimental designs for phase I and phase I/II dose-finding studies. *Brit. J. Canc.* 94 (5), 609–613.
- [Oron(2004)] Oron, A., 2004. Summary letter to M.J. Souter, MD. Consulting summary report, Statistical Consulting Unit, Department of Statistics, University of Washington, Seattle, WA, USA, available upon request from author or department.
- [Oron(2005)] Oron, A., 2005. The up-and-down experimental method: Stochastic properties and estimators. Dependent data preliminary exam report, Statistics Department, University of Washington, Seattle, WA, USA, revised version, March 2005; available from the author.

- [Oron(2006)] Oron, A., 2006. Stationary and convergence properties of 'up-and-down' methods. Technical Report 506, Department of Statistics, University of Washington, Seattle, WA, USA.
- [Oron and Hoff(In Preparation)] Oron, A., Hoff, P., In Preparation. Nonparametric estimation of up-and-down experiments, in preparation.
- [Pace and Stylianou(2007)] Pace, N., Stylianou, M., 2007. Advances in and limitations of up-and-down methodology: A prcis of clinical use, study design, and dose estimation in anesthesia research. *Anesthesiology* 107 (1), 144–152.
- [Palmer(2002)] Palmer, C., 2002. Ethics, data-dependent designs, and the strategy of clinical trials: time to start learning-as-we-go? *Stat. Meth. Med. Res.* 11 (5), 381–402.
- [Paul and Fisher(2001)] Paul, M., Fisher, D., 2001. Are estimates of MAC reliable? *Anesthesiology* 95 (6), 1362–1370.
- [Picard and Tramér(2000)] Picard, P., Tramér, M., 2000. Prevention of pain on injection with propofol: A quantitative systematic review. *Anesth. Analg.* 90.
- [Pisters et al.(2004)]Pisters, Patel, Prieto, Thall, Lewis, Feig, Hunt, Yasko, Lin, Jacobson, Burgess, Pollock, Z  
Pisters, P. W., Patel, S. R., Prieto, V. G., Thall, P. F., Lewis, V. O., Feig, B. W., Hunt, K. K., Yasko, A. W., Lin, P. P., Jacobson, M. G., Burgess, M. A., Pollock, R. E., Zagars, G. K., Benjamin, R. S., Ballo, M. T., 2004. Phase I Trial of Preoperative Doxorubicin-Based Concurrent Chemoradiation and Surgical Resection for Localized Extremity and Body Wall Soft Tissue Sarcomas. *J Clin Oncol* 22 (16), 3375–3380.  
URL <http://www.jco.org/cgi/content/abstract/22/16/3375>
- [Pollak et al.(2006)]Pollak, Palazotto, and Nicholas] Pollak, R., Palazotto, A., Nicholas, T., 2006. A simulation-based investigation of the staircase method for fatigue strength testing. *Mech. Mat.* 38 (12), 1170–1181.
- [Pollard et al.(2002)]Pollard, Makky, McFadzean, Ainsworth, Goobie, and Montgomery] Pollard, R., Makky, S., McFadzean, J., Ainsworth, L., Goobie, S., Montgomery, C., 2002. An admixture of 3 mg/kg of propofol and 3 mg/kg of thiopentone reduces pain on injection in pediatric anesthesia. *CAN. J. ANAESTH.* 49.
- [Potter(2002)] Potter, D. M., 2002. Adaptive dose finding for phase I clinical trials of drugs used for chemotherapy of cancer. *Stat. Med.* 21, 1805–1823.
- [R Development Core Team(2005)] R Development Core Team, 2005. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria, ISBN 3-900051-07-0.  
URL <http://www.R-project.org>

- [Ramsay(1998)] Ramsay, J., 1998. Estimating smooth monotone functions. *J. Royal. Stat. Soc. B* 25, 365–375.
- [Reiner et al.(1999)Reiner, Paoletti, and OQuigley] Reiner, E., Paoletti, X., OQuigley, J., 1999. Operating characteristics of the standard phase I clinical trial design. *Comp. Stat. Dat. Anal.* 30, 303–315.
- [Robbins and Monro(1951)] Robbins, H., Monro, S., 1951. A stochastic approximation method. *Ann. Math. Statistics* 22, 400–407.
- [Robertson et al.(1988)Robertson, Wright, and Dykstra] Robertson, T., Wright, F. T., Dykstra, R. L., 1988. Order restricted statistical inference. *Wiley Series in Probability and Mathematical Statistics: Probability and Mathematical Statistics*. John Wiley & Sons, Chichester.
- [Rosenberger et al.(1997)Rosenberger, Flournoy, and Durham] Rosenberger, W., Flournoy, N., Durham, S. D., 1997. Asymptotic normality of maximum likelihood estimators from multiparameter response-driven designs. *J. Statist. Plann. Inference* 60 (1), 69–76.
- [Rosenberger and Grill(1997)] Rosenberger, W., Grill, S., 1997. A sequential design for psychophysical experiments: An application to estimating timing of sensory events. *Stat. Med.* 16, 2245–2260.
- [Rosenberger and Haines(2002)] Rosenberger, W., Haines, L., 2002. Competing designs for phase I clinical trials: a review. *Stat. Med.* 21, 2757–2770.
- [Scherrer et al.(2003)Scherrer, Wiskott, Coto-Hunziker, and Belser] Scherrer, S., Wiskott, A., Coto-Hunziker, V., Belser, U., 2003. Monotonic flexure and fatigue strength of composites for provisional and definitive restoration. *J. Prosth. Dentist.* 89 (6), 579–588.
- [Shen and O’Quigley(1996)] Shen, L., O’Quigley, J., 1996. Consistency of continual reassessment method under model misspecification. *Biometrika* 83 (2), 395–405.  
URL  
<http://biomet.oxfordjournals.org/cgi/content/abstract/83/2/395>
- [Storer(1989)] Storer, B. E., 1989. Design and analysis of phase I clinical trials. *Biometrics* 45 (3), 925–937.
- [Storer(1993)] Storer, B. E., 1993. Small-sample confidence sets for the MTD in a phase I clinical trial. *Biometrics* 49 (4), 1117–1125.
- [Storer(2001)] Storer, B. E., 2001. An evaluation of phase I clinical trial designs in the continuous dose-response setting. *Stat. Med.* 20, 2399–2408.

- [Stylianou and Flournoy(2002)] Stylianou, M., Flournoy, N., 2002. Dose finding using the biased coin up-and-down design and isotonic regression. *Biometrics* 58 (1), 171–177.
- [Stylianou and Follmann(2004)] Stylianou, M., Follmann, D., 2004. The accelerated biased coin up-and-down design in phase I trials. *J. Biopharm. Stat.* 14 (1), 249–260.
- [Stylianou et al.(2003)Stylianou, Proschan, and Flournoy] Stylianou, M., Proschan, M., Flournoy, N., 2003. Estimating the probability of toxicity at the target dose following an up-and-down design. *Stat. Med.* 22 (4), 535–543.
- [Sunderam et al.(2004)Sunderam, Patra, Julli, and Warne] Sunderam, R., Patra, R., Julli, M., Warne, M., 2004. Use of the up-and-down acute toxicity test procedure to generate LC50 data for fish. *Bull. Environ. Contam. Toxic.* 72 (5), 873–880.
- ]SvenssonDeMare99 Svensson, T., de Mar{’e, J., 1999. Random features of the fatigue limit. *Extremes* 2.
- [Thall and Lee(2003)] Thall, P. F., Lee, S.-J., 2003. Practical model-based dose-finding in phase i clinical trials: Methods based on toxicity. *International Journal of Gynecological Cancer* 13 (3), 251–261.  
URL  
<http://www.blackwell-synergy.com/doi/abs/10.1046/j.1525-1438.2003.13202.x>
- [Treutwein(1995)] Treutwein, B., 1995. Minireview: adaptive psychophysical procedures. *Vision Res.* 35, 2503–2522.
- [Tsutakawa(1967a)] Tsutakawa, R., 1967a. Asymptotic properties of block up-and-down method in bio-assay. *Ann. Math. Statist.* 38 (6), 842–856.
- [Tsutakawa(1967b)] Tsutakawa, R., 1967b. Random walk design in bio-assay. *J. Amer. Statist. Assoc.* 62, 842–856.
- [Vinik et al.(1999)Vinik, Bradley, and Kissin] Vinik, H., Bradley, E., Kissin, I., 1999. Isobolographic analysis of propofol-thiopental hypnotic interaction in surgical patients. *Anesth. Analg.* 88 (5), 667–670.
- [von Bekesy(1947)] von Bekesy, G., 1947. A new audiometer. *Acta Oto.Laryn.* 35, 411–422.
- [Watson and Pelli(1979)] Watson, A. B., Pelli, D. G., 1979. The QUEST staircase procedure. *Appl. Vis. News.* 14, 6–7.
- [Watson and Pelli(1983)] Watson, A. B., Pelli, D. G., 1983. QUEST: a bayesian adaptive psychometric method. *Percept. Psychophys.* 33 (2), 113–120.

- [Weiss(1994)] Weiss, G. H., 1994. Aspects and applications of the random walk. Random Materials and Processes. North-Holland Publishing Co., Amsterdam.
- [Wetherill(1963)] Wetherill, G. B., 1963. Sequential estimation of quantal response curves. *J. Royal. Stat. Soc. B* 25, 1–48.
- [Wetherill et al.(1966)Wetherill, Chen, and Vasudeva] Wetherill, G. B., Chen, H., Vasudeva, R. B., 1966. Sequential estimation of quantal response curves: A new method of estimation. *Biometrika* 53, 439–454.
- [Wetherill and Levitt(1966)] Wetherill, G. B., Levitt, H., 1966. Sequential estimation of on a psychometric function. *Brit. J. Math. Stat. Psych.* 18, 1–10.
- [Williams(1986)] Williams, D., 1986. Interval estimation of the median lethal dose. *Biometrics* 42 (3), 641–645.
- [Xu et al.(2007)Xu, Tighiouart, and Rogatko] Xu, Z., Tighiouart, M., Rogatko, A., 2007. EWOC 2.0: Interactive software for dose escalation in cancer phase I clinical trials. *DRUG INF. J.* 41, 221–228.
- [Yuan and Chappell(2004)] Yuan, Z., Chappell, R., 2004. Isotonic designs for phase I cancer clinical trials with multiple risk groups. *Clin. Trials* 1 (6), 499–508.  
URL <http://ctj.sagepub.com/cgi/content/abstract/1/6/499>
- [Zhang and Kececioglu(1998)] Zhang, J., Kececioglu, D., 1998. New approaches to determine the endurance strength distribution.